# EXCALIBUR<sup>TM</sup> 10 EC

(ALPHA-CYPERMETHRIN 100 G/L EC)

**EXCALIBUR™ 10 EC** is a registered trademark of Agro Tico, Costa Rica **EXCALIBUR™ 10 EC** is manufactured by Chimac-Agriphar, Belgium



CHIMAC-AGRIPHAR S.A. Rue de Renory, 26 B-4102 OUGREE - BELGIUM Tel: +32-4-385.97.11 - Fax: +32-4-385.97.49

Tel: +32-4-385.97.11 - Fax: +32-4-385.97.49 Website: agriphar.com or agriphar.be

### **USE**

EXCALIBUR 10 EC is an emulsifiable concentrate insecticide containing alpha-cypermethrin, a synthetic pyrethroid which has contact, stomach, respiratory action and repellent effect. This broad-spectrum insecticide combine rapid action, good persistence and high activity on a wide range of sucking and chewing insect pests, especially Lepidoptera species, in vegetables, fruit trees, field crops and ornamental plants.

### **METHOD OF APPLICATION**

EXCALIBUR 10 EC has to be diluted with water for application by ground or air equipment.

<u>PREPARATION OF SPRAY MIXTURE</u>: Fill spray tank 1/2 full of water. Add the product directly to spray tank and mix thoroughly using mechanical or hydraulic means (complete the filling with water). Use immediately after mixing. The emulsification is immediate.

<u>APPLICATION</u>: Apply 200 to 400 litres of mixture per ha. Apply at the recommended rates when insects first appear. Use sufficient water with fine spray droplets at low volume to obtain thorough, uniform coverage.

<u>COMPATIBILITY</u>: Avoid mixtures of several materials and very concentrated spray mixtures. Incompatible with highly alkaline solutions.



CHIMAC-AGRIPHAR S.A. Rue de Renory, 26 B-4102 OUGREE - BELGIUM

Tel: +32-4-385.97.11 - Fax: +32-4-385.97.49

### **RATES OF APPLICATION**

Unless otherwise stated, the recommended rate dose is within 7 to 30 g of active ingredient per ha, depending on the conditions of application, pest species and type of crop. Follow the rates of application for each crop listed in the table on the following page.

CROPS	INSECT PESTS	RATES OF APPLICATION
Vines	Grapeberry moths Green leaf hopper Vine tortrix	100-200 ml/ha 200-275 ml/ha 100-200 ml/ha
Citrus	Aspidiotus scale Citrus aphid Flower moth, Mealy bugs	20 ml/hl water 20 ml/hl water 30-55 ml/hl water
Pome and stone fruit	Apple-aphid, Fruit moths, Pear sucker, Summer fruit tortirix moth	10-20 ml/hl water
Cotton	Armyworms Leafminer Loopers, Pink Bollworms	70-150 ml/ha 175-350 ml/ha 175-250 ml/ha
Ornamentals and flowers	Aphids, leafminers	200-300 ml/ha
Cereals and maize	Cereal bugs Maize stalk borer	70-150 ml/ha 100-250 ml/ha
Vegetables	Cereal bugs Maize stalk borer	70-150 ml/ha 100-250 ml/ha
Rice	Rice leaf folder, leafhopper	100-300 ml/ha
Soyabeans	Carrot and cabbage looper	150-350 ml/ha
Tobacco	Tobacco whitefly, flea beetles	175-275 ml/ha
Теа	Tea tortrix	10-20 ml/hl water
Coffee	White coffee leafminer	30-50 ml/ha
Oil palm	Sibine	8-35 ml/hl water



CHIMAC-AGRIPHAR S.A. Rue de Renory, 26 B-4102 OUGREE - BELGIUM

Tel: +32-4-385.97.11 - Fax: +32-4-385.97.49

### RECOMMENDATIONS

Further treatments after 10-14 days have only to be decided in function of the level of infestation. Good crop penetration and coverage is required to achieve satisfactory results, particularly when the pests are in the lower half of the crop.

### PRE-HARVEST INTEVAL

To avoid excessive residue at harvest, do not exceed the maximum dosage per ha, or the time restriction between last application and harvest for each crop: vegetables, vines, fruit trees: 14 days; cotton: 21 days; cereals: 7 days. Follow the local requirements.

### **WARNING**:

Dangerous to bees. Do not apply at flowering stage except for cereals.

### **PRECAUTIONS OF USE**

- Keep out of reach of children, away from food, drink or feed.
- Wear suitable protective gloves, face-shield and coverall.
- When using, do not eat, drink or smoke.
- Do not breathe spray mist.
- Wash any contamination from skin or eyes immediately.
- Wash hands and exposed skin before eating, drinking or smoking, before meals and after works.
- If you feel unwell, seek medical advice (show label where possible).

### **STORAGE AND DISPOSAL**

- Store in the original container, tightly closed, in a cool, well ventilated local, protected from light and heat.
- **Toxic to fish**: do not contaminated ponds, waterways or ditches with chemical or used container.
- Do not re-use this container for any other purpose. Destroy rinsed container by puncturing and crushing and dispose of safely.



CHIMAC-AGRIPHAR S.A. Rue de Renory, 26 B-4102 OUGREE - BELGIUM

Tel: +32-4-385.97.11 - Fax: +32-4-385.97.49

### **FIRST AIDS**

In case of contact with skin, remove all contaminated clothing and wash the patient thoroughly with plenty of water and soap. In case of eye contact, flush eyes with copious amounts of water for at least 15 minutes. If product has been swallowed, do not induce vomiting; call immediately a physician. Keep the patient at absolute rest.

### **NOTE TO PHYSICAN**

If a large quantity of the compound has been swallowed, gastric washing should be performed using a 5 % sodium bicarbonate solution, followed by powdered activated charcoal. Use artificial respiration, if required. Treat symptomatically.

### **CONDITIONS OF SUPPLY**

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label, only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. In no case shall seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.



CHIMAC-AGRIPHAR S.A. Rue de Renory, 26 B-4102 OUGREE - BELGIUM

Tel: +32-4-385.97.11 - Fax: +32-4-385.97.49